



Serial No.: 10/816,720
Exhibit: 2

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed



for



Limits

Preview/Index

History

Clipboard

Details

Display Abstract



Show

20



Sort by



Send to



All: 1

Review: 0



About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: Exp Cell Res. 1995 Oct;220(2):363-73.

Related Articles, Links

ELSEVIER
FULL-TEXT ARTICLE**Biochemical analysis of murine Wnt proteins reveals both shared and distinct properties.****Burrus LW, McMahon AP.**

Department of Molecular and Cellular Biology, Harvard University, Cambridge, Massachusetts 02138, USA.

The murine Wnt family of proteins consists of at least 12 members that possess significant amino acid homology. Current evidence suggests that these proteins are secreted cell-signaling molecules which are likely to have multiple roles during both embryonic development and oncogenesis. Although the biochemical properties of Wnt-1 have been thoroughly examined, less is known about the characteristics of other Wnt family members. We have compared the properties of six murine Wnt proteins (Wnt-1, Wnt-3a, Wnt-5a, Wnt-5b, Wnt-6, and Wnt-7b) transiently expressed in COS cells. All members enter the endoplasmic reticulum (ER) and are glycosylated. However, all six Wnt proteins are primarily retained in the ER in association with BiP, a resident ER protein that binds to improperly folded proteins and prevents their secretion and/or promotes proper folding. Although all Wnt family members examined are similarly processed, one notable difference was identified. Whereas addition of suramin to COS cell cultures significantly increases the levels of all six Wnts in the medium, the addition of heparin only influences the levels of Wnt-1, Wnt-6, and Wnt-7b.

PMID: 7556445 [PubMed - indexed for MEDLINE]

Display Abstract



Show

20



Sort by



Send to

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jun 27 2005 04:57:20